,

On the numerical solution of the Klein-Gordon equation by Exponential B-spline collocation method

OZLEM ERSOY HEPSON[a], ALPER KORKMAZ[b,*], IDIRIS DAG[c]

[a]DEPARTMENT OF MATHEMATICS & COMPUTER, ESKIŞEHIR OSMANGAZI UNIVERSITY, 26480, ESKIŞEHIR, TURKEY.

[b,*]DEPARTMENT OF MATHEMATICS, ÇANKIRI KARATEKIN UNIVERSITY, 18100, ÇANKIRI, TURKEY.

[c]DEPARTMENT OF COMPUTER ENGINEERING, ESKIŞEHIR OSMANGAZI UNIVERSITY, 26480, ESKIŞEHIR, TURKEY.

arXiv:1610.05649v1 [math.GM] 15 Oct 2016

ABSTRACT. In the present study, we solve initial boundary value problem construted on nonlinear Klein-Gordon equation. The collocation method on exponential cubic B-spline functions forming a set of basis for the functions defined in the same interval is set up for the numerical approach. The efficiency and validity of the proposed method are determined by computing the error between the numerical and the analytical solutions. The preservation of conserved quantities is also a good indicator of validity of the method.

1. INTRODUCTION

The Nonlinear Klein-Gordon Equation (KGE) with cubic nonlinear term is given in a generalized form

$$\frac{\partial^2 u}{\partial t^2} - \frac{\partial^2 u}{\partial x^2} + P'(u) = 0 \tag{1.1}$$

where $u = u(x,t)$ and $P(u)$ is reasonable nonlinear function that is chosen as the potential energy in many cases [1]. The choice of $P'(u)$ can generate various equations such as Sine-Gordon, Landau-Ginzburg-Higgs, ϕ^4 or reaction Duffing equations [2], arising to model many physical phenomena from quantum mechanics to wave motion. The origin of the KGE iss discussed in the Kragh's essay [3]. He mentions in his essay that some earlier types of the KGE was connected with the general relativity theory by siginificient physicists covering Klein, Fock and Schrödinger. The KGE is absolutely one of the most important equation that the nonlinear Schrödinger equation can be derived from this equation in the canonical form [4]. Galehouse [5] derived the Klein-Gordon equation geometrically using appropriate gauge transformations.

The scattering theory for the more general KGE is proved by Schechter [6]. The same theory for the KGE is developed by following classical procedure of an equaivalent equation containing first order time in a particular finite energy normed Hilbert space [7]. Weder gave the proofs of the existence and completeness of the wave operators, the intertwining relations, and the invariance principle in that study [7]. Spectral and scattering theory of the KGE are derived by implementing related eignfunction expansions in a particular field case to give strong results like in the Schrödinger case [8]. Motion of a KGE in an external field is examined by Tsukanov [9]. The linear response approximation is used to calculate the drift velocity of a kink of the KGE in study.

Since introduced, many types of solutions describing various physical phenomena covering solitary waves and kinks have been found for the KGE. Derivations of solutions of the KGE from the solutions of the wave equation are studied by Chambers [10]. Some bound state type solutions of the KGE for various potentials are set up explicitly by Fleischer and Soff [11]. An exact formal solution for a particular nonlinear KGE is obtained by manipulating the solution of linear Klein-Gordon equation [12]. The envelope soliton and hole solutions for the nonlinear KGE are studied by transforming it to the nonlinear Schrödinger equation [13]. Some solitary wave solutions of a class of field equations for systems with polynomial self-interactions reduce to plane-wave solutions of the KGE [14]. Some bell and kink type solitary wave solutions formed by using some hyperbolic functions for the KGE are constructed with the help of the extended first kind elliptic sub-equation method [15]. Some similar solutions have been derived by Akter and Akbar using modified simple equation method [16]. Adomian implemented the decomposition method to the initial boundary value problems for the KGE [17]. The interaction of two solitons that are self-localized and weakly oscillating and damped for the particular form of the KGE are studied by Kudryavtsev [18].

Besides some analytical solutions summarized above, various numerical methods have been applied to derive solutions of the problems related to the KGE. Strauss and Vazquez [19] solved some problems constructed on the KGE by the simplest central second difference method. The equation integrated numerically by using four explicit finite

2010 *Mathematics Subject Classification.* 35K57,65L60,92E20.

Key words and phrases. *Klein-Gordon Equation, exponential cubic B-spline, collocation, wave motion.*

Corresponding author: Alper Korkmaz; alperkorkmaz7@gmail.com

difference methods resulting that the energy conserving scheme is more suitable than the others to simulate the long time behaviour of the solutions [20]. A collocation method based on thin plate splines radial basis functions is another proposed method to solve the KGE numerically [21]. The solutions of the nonhomogenous form of the KGE are approximated by multiquadric quasi-interpolation and the integrated radial basis function network schemes [22]. Classical polynomial cubic B-spline functions have also been adapted for the collocation method to derive numerical solutions of the KGE [23, 24].

In the present study, we solve some initial boundary value problems derived for the nonlinear KGE by the collocation method based on nonpolynomial B-splines, namely exponential cubic B-spline functions.

In order to solve the initial boundary value problem setup for the nonlinear KGE (1.1) numerically, we first reduce the order of the derivative in time for a particular case assuming $-P'(u) = u - u^3$

$$\begin{aligned} v_t &= u_{xx} + u - u^3 \\ u_t &= v \end{aligned} \tag{1.2}$$

To complete the usual classical mathematical statement of the problem, the initial and the boundary conditions are chosen as

$$\begin{aligned} u(x,0) &= f(x), \quad a \le x \le b \\ v(x,0) &= g(x), \quad a \le x \le b \end{aligned} \tag{1.3}$$

and

$$u_x(a,t) = u_x(b,t) = 0, v_x(a,t) = v_x(b,t) = 0 \tag{1.4}$$

in the finite problem domain $[a, b]$.

2. Exponential B-spline Collocation Method

Let κ be a uniformly distributed grids of the finite interval $[a, b]$, such as,

$$\kappa : x_m = a + mh, m = 0, 1, \dots N$$

where $h = \frac{b-a}{N}$. Then, the exponential cubic B-splines set

$$B_m(x) = \begin{cases} b_2\left((x_{m-2} - x) - \frac{1}{\rho}\sinh(\rho(x_{m-2} - x))\right) & , \quad [x_{m-2}, x_{m-1}] \\ a_1 + b_1(x_m - x) + c_1\exp(\rho(x_m - x)) + d_1\exp(-\rho(x_m - x)) & , \quad [x_{m-1}, x_m] \\ a_1 + b_1(x - x_m) + c_1\exp(\rho(x - x_m)) + d_1\exp(-\rho(x - x_m)) & , \quad [x_m, x_{m+1}] \\ b_2\left((x - x_{m+2}) - \frac{1}{\rho}\sinh(\rho(x - x_{m+2}))\right) & , \quad [x_{m+1}, x_{m+2}] \\ 0 & , \quad otherwise \end{cases} \tag{2.5}$$

where $m = -1, 0, \dots N$,

$$\begin{aligned} a_1 &= \frac{\rho h\cosh(\rho h)}{\rho h\cosh(\rho h) - \sinh(\rho h)}, \\ b_1 &= \frac{\rho}{2}\frac{\cosh(\rho h)(\cosh(\rho h) - 1) + \sinh^2(\rho h)}{(\rho h\cosh(\rho h) - \sinh(\rho h))(1 - \cosh(\rho h))}, \\ b_2 &= \frac{\rho}{2(\rho h\cosh(\rho h) - \sinh(\rho h))}, \\ c_1 &= \frac{1}{4}\frac{\exp(-\rho h)(1 - \cosh(\rho h)) + \sinh(\rho h)(\exp(-\rho h) - 1))}{(\rho h\cosh(\rho h) - \sinh(\rho h))(1 - \cosh(\rho h))}, \\ d_1 &= \frac{1}{4}\frac{\exp(\rho h)(\cosh(\rho h) - 1) + \sinh(\rho h)(\exp(\rho h) - 1))}{(\rho h\cosh(\rho h) - \sinh(\rho h))(1 - \cosh(\rho h))}, \end{aligned}$$

with real parameter ρ forms a basis for the functions defined in $[a, b]$ [25]. Each exponential cubic B-spline $B_m(x)$ has two continuous lowest derivatives defined in the interval $[x_{m-1}, x_{m+2}]$. Different from polynomial B-splines [26],this basis function set has been chosen as trial functions in various methods to solve different problems arising in different engineering fields [27–32]. The approximate solutions U and V can be written in terms of the exponential B-splines as

$$\begin{aligned} U(x,t) &= \sum_{i=-1}^{N+1}\delta_i B_i(x), \\ V(x,t) &= \sum_{i=-1}^{N+1}\varphi_i B_i(x) \end{aligned} \tag{2.6}$$

where δ_i and φ_i are time dependent parameters to be determined from the complementary conditions. The nodal values of U, V and their lowest two derivatives at the nodes can be found from (2.6) as

$$\begin{aligned} U_i &= U(x_i,t) = \frac{s-\rho h}{2(\rho hc-s)}\delta_{i-1} + \delta_i + \frac{s-\rho h}{2(\rho hc-s)}\delta_{i+1}, \\ U_i' &= U'(x_i,t) = \frac{\rho(1-c)}{2(\rho hc-s)}\delta_{i-1} + \frac{\rho(c-1)}{2(\rho hc-s)}\delta_{i+1} \\ U_i'' &= U''(x_i,t) = \frac{\rho^2 s}{2(\rho hc-s)}\delta_{i-1} - \frac{\rho^2 s}{\rho hc-s}\delta_i + \frac{\rho^2 s}{2(\rho hc-s)}\delta_{i+1}. \end{aligned} \tag{2.7}$$

$$\begin{aligned} V_i &= V(x_i,t) = \frac{s-\rho h}{2(\rho hc-s)}\phi_{i-1} + \phi_i + \frac{s-\rho h}{2(\rho hc-s)}\phi_{i+1}, \\ V_i' &= V'(x_i,t) = \frac{\rho(1-c)}{2(\rho hc-s)}\phi_{i-1} + \frac{\rho(c-1)}{2(\rho hc-s)}\phi_{i+1} \\ V_i'' &= V''(x_i,t) = \frac{\rho^2 s}{2(\rho hc-s)}\phi_{i-1} - \frac{\rho^2 s}{\rho hc-s}\phi_i + \frac{\rho^2 s}{2(\rho hc-s)}\phi_{i+1}. \end{aligned} \tag{2.8}$$

where $c = \cosh \rho h$ and $s = \sinh \rho h$.
The time discretization of the system (1.2) by usual forward finite difference and the Crank-Nicolson method leads to

$$\begin{aligned} \frac{V^{n+1}-V^n}{\Delta t} - \frac{U_{xx}^{n+1}+U_{xx}^n}{2} - \frac{U^{n+1}+U^n}{2} + \frac{(U^3)^{n+1}+(U^3)^n}{2} = 0 \\ \frac{U^{n+1}-U^n}{\Delta t} - \frac{V^{n+1}+V^n}{2} = 0 \end{aligned} \tag{2.9}$$

where $U^{n+1} = U(x,(n+1)\Delta t)$ represents the solution at the $(n+1)$.th time level. Linearizing the term $(U^3)^{n+1}$ in (2.9) as

$$(U^3)^{n+1} = 3U^{n+1}(U^2)^n - 2(U^3)^n \tag{2.10}$$

yields the time-integrated system:

$$\begin{aligned} \frac{V^{n+1}-V^n}{\Delta t} - \frac{U_{xx}^{n+1}+U_{xx}^n}{2} - \frac{U^{n+1}+U^n}{2} + \frac{3U^{n+1}(U^2)^n-(U^3)^n}{2} = 0 \\ \frac{U^{n+1}-U^n}{\Delta t} - \frac{V^{n+1}+V^n}{2} = 0 \end{aligned} \tag{2.11}$$

Approximating to U and V by using the nodal values (2.7) and (2.8) gives

$$\begin{aligned} &\nu_{m1}\delta_{m-1}^{n+1} + \nu_{m2}\phi_{m-1}^{n+1} + \nu_{m3}\delta_m^{n+1} + \nu_{m4}\phi_m^{n+1} + \nu_{m1}\delta_{m+1}^{n+1} + \nu_{m2}\phi_{m+1}^{n+1} \\ = \;& \nu_{m5}\delta_{m-1}^{n} + \nu_{m2}\phi_{m-1}^{n} + \nu_{m6}\delta_m^{n} + \nu_{m4}\phi_m^{n} + \nu_{m5}\delta_{m+1}^{n} + \nu_{m2}\phi_{m+1}^{n} \end{aligned} \tag{2.12}$$

$$\begin{aligned} &\nu_{m2}\delta_{m-1}^{n+1} + \nu_{m7}\phi_{m-1}^{n+1} + \nu_{m4}\delta_m^{n+1} + \nu_{m8}\phi_m^{n+1} + \nu_{m2}\delta_{m+1}^{n+1} + \nu_{m7}\phi_{m+1}^{n+1} \\ = \;& \nu_{m2}\delta_{m-1}^{n} - \nu_{m7}\phi_{m-1}^{n} + \nu_{m4}\delta_m^{n} - \nu_{m8}\phi_m^{n} + \nu_{m2}\delta_{m+1}^{n} - \nu_{m7}\phi_{m+1}^{n} \end{aligned} \tag{2.13}$$

The coefficients of iteration system (2.12)-(2.13) are

$$\begin{aligned} \nu_{m1} &= \left(3K^2-1\right)\alpha_1 - \gamma_1 \\ \nu_{m2} &= \frac{2}{\Delta t}\alpha_1 \\ \nu_{m3} &= \left(3K^2-1\right)\alpha_2 - \gamma_2 \\ \nu_{m4} &= \frac{2}{\Delta t}\alpha_2 \\ \nu_{m5} &= \left(1+K^2\right)\alpha_1 + \gamma_1 \\ \nu_{m6} &= \left(1+K^2\right)\alpha_2 + \gamma_2 \\ \nu_{m7} &= -\alpha_1 \\ \nu_{m8} &= -\alpha_2 \end{aligned}$$

where

$$K = \alpha_1\delta_{i-1} + \alpha_2\delta_i + \alpha_1\delta_{i+1}$$

and

$$\begin{aligned} \alpha_1 &= \frac{s-ph}{2(phc-s)},\ \alpha_2 = 1, \\ \gamma_1 &= \frac{p^2 s}{2(phc-s)},\ \gamma_2 = -\frac{p^2 s}{phc-s} \end{aligned}$$

The system system (2.12)-(2.13) can be written in the matrix form

$$\mathbf{A}\mathbf{x}^{n+1} = \mathbf{B}\mathbf{x}^{n} \tag{2.14}$$

where

$$\mathbf{A} = \begin{bmatrix} \nu_{m1} & \nu_{m2} & \nu_{m3} & \nu_{m4} & \nu_{m1} & \nu_{m2} & & & & \\ \nu_{m2} & \nu_{m7} & \nu_{m4} & \nu_{m8} & \nu_{m2} & \nu_{m7} & & & & \\ & & \nu_{m1} & \nu_{m2} & \nu_{m3} & \nu_{m4} & \nu_{m1} & \nu_{m2} & & \\ & & \nu_{m2} & \nu_{m7} & \nu_{m4} & \nu_{m8} & \nu_{m2} & \nu_{m7} & & \\ & & & \ddots & \ddots & \ddots & \ddots & \ddots & \ddots & \\ & & & & \nu_{m1} & \nu_{m2} & \nu_{m3} & \nu_{m4} & \nu_{m1} & \nu_{m2} \\ & & & & \nu_{m2} & \nu_{m7} & \nu_{m4} & \nu_{m8} & \nu_{m2} & \nu_{m7} \end{bmatrix}$$

and

$$\mathbf{B} = \begin{bmatrix} \nu_{m5} & \nu_{m2} & \nu_{m6} & \nu_{m4} & \nu_{m5} & \nu_{m2} & & & & \\ \nu_{m2} & -\nu_{m7} & \nu_{m4} & -\nu_{m8} & \nu_{m2} & -\nu_{m7} & & & & \\ & & \nu_{m5} & \nu_{m2} & \nu_{m6} & \nu_{m4} & \nu_{m5} & \nu_{m2} & & \\ & & \nu_{m2} & -\nu_{m7} & \nu_{m4} & -\nu_{m8} & \nu_{m2} & -\nu_{m7} & & \\ & & & \ddots & \ddots & \ddots & \ddots & \ddots & \ddots & \\ & & & & \nu_{m5} & \nu_{m2} & \nu_{m6} & \nu_{m4} & \nu_{m5} & \nu_{m2} \\ & & & & \nu_{m2} & -\nu_{m7} & \nu_{m4} & -\nu_{m8} & \nu_{m2} & -\nu_{m7} \end{bmatrix}$$

This system consists of $2N+2$ linear equations with $2N+6$ unknown parameters $\mathbf{x}^{n+1} = (\delta_{-1}^{n+1}, \phi_{-1}^{n+1}, \delta_0^{n+1}, \phi_0^{n+1} \ldots, \delta_{N+1}^{n+1}, \phi_{N+1}^{n+1}$
A unique solution of the system can be obtained by imposing the boundary conditions $U_x(a,t) = 0, U_x(b,t) = 0, V_x(a,t) = 0, V_x(b,t) = 0$ to have the following the equations:

$$\delta_{-1} = \delta_1,\ \phi_{-1} = \phi_1,\ \delta_{N-1} = \delta_{N+1},\ \phi_{N-1} = \phi_{N+1}$$

Elimination of the parameters $\delta_{-1}, \phi_{-1}, \delta_{N+1}, \phi_{N+1}$, using the Eqs,(2.12) and (2.13) from the the system gives a solvable system of $2N+2$ linear equation including $2N+2$ unknown parameters. Placing solution parameters when is computed from the system via a variant of the Thomas algorithm gives the approximate solution over the subregion $[x_i, x_{i+1}]$. We need the initial parameter vectors $d_1 = (\delta_{-1}, \delta_0, ..\delta_N, \delta_{N+1})$, $d_2 = (\phi_{-1}, \phi_0, ..\phi_N, \phi_{N+1})$ to start the iteration process for the system of equation: To do that, the following requirements help to determine initial parameters:

$U_x(a,0) = 0 = \delta_{-1}^0 - \delta_1^0$,
$U_x(x_i,0) = \delta_{i-1}^0 - \delta_{i+1}^0, i = 1, ..., N-1$
$U_x(b,0) = 0 = \delta_{N-1}^0 - \delta_{N+1}^0$,
$V_x(a,0) = 0 = \phi_{-1}^0 - \phi_1^0$
$V_x(x_i,0) = \phi_{i-1}^0 - \phi_{i+1}^0, i = 1, ..., N-1$
$V_x(b,0) = 0 = \phi_{N-1}^0 - \phi_{N+1}^0$

3. Numerical Example

We illustrate the performance of the proposed method in terms of the accuracy. The absolute error between the obtained numerical solution and the exact solution is computed for different values of time step length Δt and step size h. The discrete maximum norm

$$L_\infty = |w - W|_\infty = \max_j |w_j - W_j|$$

is computed where w and W represent the analytical and numerical solutions, respectively.
The conservation laws defined as [33]

$$E = \tfrac{1}{2}\int_a^b \left[V^2 + U_x^2 - U^2 + \tfrac{1}{2}U^4\right] dx$$
$$P = \tfrac{1}{2}\int_a^b V U_x dx$$

are also computed for each time step to observe if they are preserved.
The KGE equation has an analytical solution representing a travelling wave of the form

$$u(x,t) = \tanh\left(\frac{(x-ct)}{\sqrt{2(1-c^2)}}\right) \tag{3.15}$$

where c is the velocity of wave [33]. The initial condition is generated by substituting $t = 0$ into the analytical solution. In this test problem, the parameters are chosen as $\Delta t = 0.005, h = 0.02$, $c = 0.5$ and various values of the exponential B-spline parameter ρ in the interval $[a,b] = [-30, 30]$ for $t \in [0, 30]$. The maximum error comparison at the simulation

terminating time $t = 30$ shows that the results are better in two decimal digits when $\rho = 0.0000589$, Fig 1(a)-1(b). The



(a) $\rho = 1$ (b) $\rho = 0.0000589$

FIGURE 1. Comparison of error norms L_∞ for various values of ρ

maximum errors are tabulated for various values of space and time step lengths and the exponential B-spline parameter ρ at time $t = 10$ in Table 1. A comparison with Boz's work, based on polynomial B-spline collocation, shows that the exponential B-spline generates the results one decimal digit better even when the same parameters are used.

Table 1: $L_\infty \times 10^3$ error norms for $c = 0.5$, $t = 10$, $-30 \le x \le 30$

Time		Present ($\rho = 1$)	Present (various ρ)	Ref: [34]
h	Δt			
0.2	0.05	17.5000	$3.6155(\rho = 0.0000012)$	11.0155
0.1	0.02	4.5179	$0.2525(\rho = 0.0000053)$	2.8275
0.05	0.01	1.1333	$0.0583(\rho = 0.0000151)$	0.6611
0.02	0.005	0.1790	$0.0076(\rho = 0.0000589)$	0.0466

The analytical values of the conservation laws initially are calculated as $E = -13.91133798$ and $P = -0.5443310539$. The relative changes of the conservation laws defined as

$$C_P(t) = \left| \frac{P_t - P_0}{P_0} \right| \qquad C_E(t) = \left| \frac{E_t - E_0}{E_0} \right| \tag{3.16}$$

where $C_P(t)$ and $C_E(t)$ are the relative changes of conservation laws at the time t with respect to the reference initial values E_0 and P_0. We record the computed values of relative changes in Table 2. Even when the results obtained various values when $\rho = 1$ can be recovered by changing it, Table 1, a significant recovery is not observed in the relative changes of conservation laws, Table 2.

Table 2: Relative changes of conservative laws for various values of ρ

Time		$(\rho = 1)$	$(\rho = 1)$	(various ρ)	(various ρ)
h	Δt	$C_P(10)$	$C_E(10)$	$C_P(10)$	$C_E(10)$
0.2	0.05	3.9512×10^{-5}	1.3011×10^{-6}	$3.6516 \times 10^{-5}(\rho = 0.0000012)$	2.2376×10^{-7}
0.1	0.02	2.4573×10^{-6}	8.6386×10^{-8}	$2.3098 \times 10^{-6}(\rho = 0.0000053)$	2.5001×10^{-8}
0.05	0.01	2.9692×10^{-7}	2.6278×10^{-9}	$2.8803 \times 10^{-7}(\rho = 0.0000151)$	4.5722×10^{-9}
0.02	0.005	3.6163×10^{-8}	4.9621×10^{-10}	$3.5804 \times 10^{-8}(\rho = 0.0000589)$	6.8026×10^{-10}

4. CONCLUSION

In the study, we derive a collocation method based on exponential B-spline functions combined with strong stable Crank-Nicolson implicit method. In order to check the validity and accuracy of the method, we solve an initial boundary value problem constructed with nonlinear Klein-Gordon equation. The accuracy of the method is measured by computing the maximum error norm. Comparison with the results obtained by polynomial B-splines in [34] shows that the proposed method is capable and generates more accurate results. The relative changes of the lowest two conservation laws are also good indicators of good agreement of the numerical results with analytical ones. Different from polynomial B-splines, the change of parameter in the exponential B-splines provides recovery in the results.

Acknowledgements: A brief part of this study was orally presented in *the International Conference on Analysis and its Applications-ICAA2016,Kırşehir, Turkey, 2016.*

REFERENCES

[1] Whitham G.B., *Linear and Nonlinear Waves*, John Wiley & Sons Inc. New York, 1999.

[2] Kim, J. J., & Hong, W. P. (2004). *New solitary-wave solutions for the generalized reaction Duffing model and their dynamics.* Zeitschrift für Naturforschung A, **59**(11), 721-728.

[3] Kragh, H. (1984). *Equation with the many fathers. The Klein-Gordon equation in 1926.* American Journal of Physics, **52**(11), 1024-1033.

[4] Ablowitz M., *Nonlinear Dispersive Waves*, Cambridge Press, Cambridge, 2011.

[5] Galehouse, D. C. (1981). *Geometrical derivation of the Klein-Gordon equation.* International Journal of Theoretical Physics, **20**(6), 457-479.

[6] Schechter, M. (1976). *The Klein-Gordon equation and scattering theory.* Annals of Physics, **101**(2), 601-609.

[7] Weder, R. A. (1978). *Scattering theory for the Klein-Gordon equation.* Journal of Functional Analysis, **27**(1), 100-117.

[8] Lundberg, L. E. (1973). *Spectral and scattering theory for the Klein-Gordon equation.* Communications in Mathematical Physics, **31**(3), 243-257.

[9] Tsukanov, V. D. (1990). *Motion of a Klein-Gordon kink in an external field.* Theoretical and Mathematical Physics, **84**(3), 930-933.

[10] Chambers, L. G. (1966). *Derivation of solutions of the Klein-Gordon equation from solutions of the wave equation.* Proceedings of the Edinburgh Mathematical Society (Series 2), **15**(02), 125-129.

[11] Fleischer, W., & Soff, G. (1984). *Bound state solutions of the Klein-Gordon equation for strong potentials.* Zeitschrift für Naturforschung A, **39**(8), 703-719.

[12] Burt, P. B., & Reid, J. L. (1976). *Exact solution to a nonlinear Klein-Gordon equation.* Journal of Mathematical Analysis and Applications, **55**(1), 43-45.

[13] Sharma, A. S., & Buti, B. (1976). *Envelope solitons and holes for sine-Gordon and non-linear Klein-Gordon equations.* Journal of Physics A: Mathematical and General, **9**(11), 1823.

[14] Burt, P. B. (1974). *Solitary waves in nonlinear field theories.* Physical Review Letters, **32**(19), 1080.

[15] Huang, D. J., & Zhang, H. Q. (2005). *The extended first kind elliptic sub-equation method and its application to the generalized reaction Duffing model.* Physics Letters A, **344**(2), 229-237.

[16] Akter, J., & Akbar, M. A. (2015). *Exact solutions to the Benney-Luke equation and the Phi-4 equations by using modified simple equation method.* Results in Physics, **5**, 125-130.

[17] Adomian, G. (1996). *Nonlinear klein-gordon equation.* Applied Mathematics Letters, **9**(3), 9-10.

[18] Kudryavtsev, A. E. (1975). *Solitonlike solutions for a Higgs scalar field.* Institute of Theoretical and Experimental Physics.

[19] Strauss, W., & Vazquez, L. (1978). *Numerical solution of a nonlinear Klein-Gordon equation.* Journal of Computational Physics, **28**(2), 271-278.

[20] Jiménez, S., & Vázquez, L. (1990). *Analysis of four numerical schemes for a nonlinear Klein-Gordon equation.* Applied Mathematics and Computation, **35**(1), 61-94.

[21] Dehghan, M., & Shokri, A. (2009).*Numerical solution of the nonlinear Klein-Gordon equation using radial basis functions.* Journal of Computational and Applied Mathematics, **230**(2), 400-410.

[22] Sarboland, M., Aminataei, A. (2015). *Numerical Solution of the Nonlinear Klein-Gordon Equation Using Multiquadric Quasi-interpolation Scheme.* Universal Journal of Applied Mathematics, **3**(3), 40-49.

[23] Zahra, W. K., Ouf, W. A., & El-Azab, M. S. (2016). *A robust uniform B-spline collocation method for solving the generalized PHI-four equation.* Applications and Applied Mathematics, **11**(1), 384-396.

[24] Rashidinia, J., Ghasemi, M., & Jalilian, R. (2010). *Numerical solution of the nonlinear Klein-Gordon equation.* Journal of Computational and Applied Mathematics, **233**(8), 1866-1878.

[25] McCartin, B.J., *Theory of exponential splines.* Journal of Approximation Theory, **66**1, 1-23, 1991.

[26] Korkmaz, A., & Dag, I. (2013). *Cubic B-spline differential quadrature methods and stability for Burgers' equation.* Engineering Computations, **30**(3), 320-344.

[27] Mohammadi, R., *Exponential B-Spline Solution of Convection-Diffusion Equations.* Applied Mathematics, **4**, 933-944, 2013.

[28] Mohammadi, R., *Exponential B-spline collocation method for numerical solution of the generalized regularized long wave equation* Chin. Phys. B, **24** 5, 050206, 2015.

[29] Ersoy, O., & Dag, I., *The exponential cubic B-spline algorithm for Korteweg-de Vries Equation* Advances in Numerical Analysis, Article ID 367056, 2015.

[30] Ersoy, O., & Dag, I. (2016). *The Exponential Cubic B-Spline Collocation Method for the Kuramoto-Sivashinsky Equation.* Filomat, **30** 3, 853-861.

[31] Dag, I., & Ersoy, O. (2016). *The exponential cubic B-spline algorithm for Fisher equation.* Chaos, Solitons & Fractals, **86**, 101-106.

[32] Ersoy, O., & Dag, I. (2015). *Numerical solutions of the reaction diffusion system by using exponential cubic B-spline collocation algorithms.* Open Physics, **13**(1).

[33] Zaki, S.I., Gardner L.R.T., Gardner G.A., 1997, *Numerical simulations of Klein-Gordon solitary wave interactions* Il Nuovo Cimento, 112B, N.7.

[34] A. Boz, *Solution of Klein Gordon Equation using B-Spline Collocation Methods.* PhD Thesis, 2006.